

Insecticide Resistance Management Strategies
Developed by the CropLife Australia Insecticide Resistance Management Review Group
 Valid as at 27 June 2013

Crop(s) : Pasture/Winter Crops
Mite : Redlegged Earth Mite (RLEM)

Guidelines:

Rotate insecticide groups.

Do not apply consecutive sprays of products from any one insecticide group

Crop Stage	Group*	Chemical Sub-group	Example chemical
Seed Treatment (or in-furrow)	4A	Neo-nicotinoids	Imidacloprid
	1B	organophosphates	dimethoate
	2B	phenylpyrazoles	fipronil
Bare Earth (Pre-emergent)	1B	organophosphates	omethoate
	3A	synthetic pyrethroids	bifenthrin
Early Season (Autumn when limited green growth)	3A	synthetic pyrethroids	alpha-cypermethrin
	1B	organophosphates	chlorpyrifos
Spring	1B	organophosphates	Omethoate
	3A	synthetic pyrethroids	gamma-cyhalothrin

*Groups are the International Resistance Action Committee Insecticide Groups based on mode of action of the insecticides - refer MoA tables.

If both autumn and spring applications are needed, alternate between synthetic pyrethroids and organophosphates

Timing of Sprays

1. Monitor Redlegged Earth Mite (RLEM) activity carefully and only treat if damage has reached threshold levels.
2. One well timed spray in Autumn or Spring will maximise effectiveness of treatment.

Placement of Sprays

1. Apply perimeter sprays where infestations are concentrated on the edge of fields.
2. Use blanket sprays where appropriate.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 27 June 2013**. All previous versions of this strategy are now invalid.
 Phone: 02 6230 6399 Email: info@croplifeaustralia.org.au Fax: 02 6230 6355 Website: www.croplifeaustralia.org.au

Crop(s) : Pasture/Winter Crops (cont.)
Mite : Redlegged Earth Mite (RLEM) (cont.)

Guidelines: (cont.)

Cultural Practices

1. Heavy grazing or cutting for hay or cultivation will reduce mite numbers.
2. Develop damage thresholds.
3. Rotate crops and pastures that are more tolerant to the pest.
4. Encourage predator survival by judicious use of insecticides.
5. Control alternative hosts such as Capeweed and Paterson's curse.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 27 June 2013**. All previous versions of this strategy are now invalid.
Phone: 02 6230 6399 Email: info@croplifeaustralia.org.au Fax: 02 6230 6355 Website: www.croplifeaustralia.org.au